# **DERMO-COSMETIC** CARE TO SUPPORT SKIN **DURING RADIOTHERAPY**

# Cicabio Restor

Protective soothing care for skin weakened by some cancer treatments

SKIN WEAKENED BY RADIOTHERAPY



## A COMPLETE ROUTINE FOR RADIOTHERAPY



### **CLEAN GENTLY**

Atoderm Intensive gel moussant\*



TREAT Cicabio Restor

### REPAIRS

Skin barrier

### **SOOTHES**

Feelings of discomfort

### **PROTECTS**

The epidermis



\*For adults and children of more than 3 years of age. Not to be used on visibly altered or oozing skin. Consult your doctor in case of severe liver or kidney disorders. In case of worsening of skin conditions, stop the applications and consult a healthcare professional

NAOS, SAS with a capital of € 43,474,650,R.C.S. Aix-en-Provence B535 236 418, 355 rue Pierre-Simon Laplace 13290 Aixen-Provence. LD-AB(0428)Sept 2021

#### BIOLOGY AT THE SERVICE OF DERMATOLOGY

Learn more about NAOS, French ecobiology company, founder of BIODERMA, on www.naos.com



# **BIODERMA** LABORATOIRE DERMATOLOGIQUE



## SKIN CARE ROUTINE

Before, during and after

Complete routine to support your skin



# RADIOTHERAPY COULD IMPACT THE SKIN

# TAKE CARE OF YOUR SKIN BEFORE, DURING AND AFTER RADIOTHERAPY



Radiotherapy limits the multiplication of cancer cells thanks to local irradiation. Even if they are more and more rare,

### RADIOTHERAPY COULD IMPACT THE SKIN and thus AFFECT YOUR QUALITY OF LIFE.




Nevertheless, with an adapted routine it is possible to prevent and **relieve** possible feelings of pain and burn.

# BEFORE YOUR FIRST RADIOTHERAPY SESSION **DURING 15 DAYS**



Start to apply once to twice a day on the concerned area a nourishing and repairing **cream** to prepare your skin 15 days before your first session.



Do not apply anything on (cream, lotion, deodorizers. perfume...).

# THE DAY OF YOUR RADIOTHERAPY SESSION REPEAT AT EACH SESSION

AFTER THE SESSION



Apply a repairing and soothing cream to limit the impact of the radiation on your skin.

AFTER YOUR LAST **SESSION** 

**DURING 15 DAYS** 



At least 2 weeks after the end of your protocol, apply on the area a repairing cream once or twice a day.